

THE LALONDE-MCDUFF CONJECTURE AND THE FUNDAMENTAL GROUP

JAREK KĘDRA

ABSTRACT. We give a simple proof of the Lalonde-McDuff conjecture for aspherical manifolds.

1. INTRODUCTION

Let G be a topological group acting on a manifold M . Let

$$M \xrightarrow{i} M_G \xrightarrow{\pi} BG$$

be the universal fibration associated to the action. It is a fundamental question to determine the cohomology $H^*(M_G)$. This cohomology is also known as the equivariant cohomology of M associated with the action of G and denoted by $H_G(M)$. A cohomology class in $H^*(M_G)$ is called a G -equivariant class of M .

It follows from the Leray-Hirsch theorem (Theorem 4D.1 in Hatcher [2]) that $H(M_G; \mathbb{Q})$ is isomorphic as an $H^*(BG; \mathbb{Q})$ -module to the tensor product of the cohomology of the base and the cohomology of the fibre if and only if the homomorphism $i^* : H^*(M_G; \mathbb{Q}) \rightarrow H^*(M; \mathbb{Q})$ induced by the inclusion of the fibre is surjective.

In the present note we make a simple observation about the above homomorphism under some hypothesis on the group action. As a consequence we prove the so called Lalonde-McDuff conjecture for aspherical manifolds (see Section 3 for details).

2. THE MAIN OBSERVATION

Let $c : M \rightarrow B\pi_1(M)$ be the map classifying the universal cover. A cohomology class $\alpha \in H^*(M)$ is called a π_1 -class if it is in the image of the induced homomorphism

$$c^* : H^*(\pi_1(M)) \rightarrow H^*(M).$$

Date: January 27, 2009.

2000 Mathematics Subject Classification. Primary 57R17; Secondary 55R10.

Key words and phrases. Hamiltonian fibration; cohomology.

Let $p \in M$ be a fixed point and let $\text{ev} : G \rightarrow M$ be the corresponding evaluation map defined by

$$\text{ev}(f) := f(p).$$

Theorem 2.1. *Suppose that the evaluation map induces the trivial homomorphism on the fundamental group. Then every π_1 -class is the image of some G -equivariant class.*

Proof. If the evaluation map induces the trivial homomorphism on the fundamental group then the connecting homomorphism

$$\partial : \pi_2(BG) \rightarrow \pi_1(M)$$

is also trivial. In fact, it is the same homomorphism as the one induced by the evaluation, after the identification $\pi_1(G) \cong \pi_2(BG)$. Since G is connected, BG is simply connected and, by the long exact sequence of homotopy groups of the universal fibration, we get the isomorphism $i_* : \pi_1(M) \rightarrow \pi_1(M_G)$. Thus the classifying map $c : M \rightarrow B\pi_1(M)$ factors through M_G which finishes the proof. $\square$

Corollary 2.2. *Let M be an aspherical manifold. If the evaluation map induces the trivial homomorphism on the fundamental group then the universal fibration $M \rightarrow M_G \rightarrow BG$ is homotopy trivial. In particular, the G -equivariant cohomology of M is isomorphic as a ring to the tensor product $H^*(BG) \otimes H^*(M)$.*

Proof. Since M is aspherical we have that $M = B\pi_1(M)$. The map $M_G \rightarrow B\pi_1(M)$ classifying the universal cover is a homotopy inverse of the inclusion of the fibre $i : M \rightarrow M_G$. This implies that the fibration is homotopy trivial. $\square$

Example 2.3. It is easy to see that the image of the homomorphism induced by the evaluation map is contained in the centre of the fundamental group of M . Hence the above results apply to manifolds whose fundamental group has trivial centre. In particular, if M is such an aspherical manifold then any bundle with fibre M over simply connected base is homotopy trivial.

Remark 2.4. The last observation also follows from the classical result of Gottlieb [1, Theorem III.2] which states that the identity component of the space of homotopy equivalences of an aspherical manifold M is itself aspherical and has the fundamental group isomorphic to the centre of $\pi_1(M)$.

3. THE LALONDE-MCDUFF CONJECTURE

Let (M, ω) be a symplectic manifold. A locally trivial bundle

$$(M, \omega) \xrightarrow{i} E \xrightarrow{\pi} B$$

is called **Hamiltonian** if its structure group is a subgroup of the group $\text{Ham}(M, \omega)$ of Hamiltonian diffeomorphisms of (M, ω) . Lalonde and McDuff conjectured in [3] that the rational cohomology of the total space is isomorphic as a vector space to the tensor product of the cohomology of the base and the cohomology of the fibre. The conjecture has been proved in many particular cases. The examples include fibrations where the fibre is a Kähler manifold, fibrations for which the structure group is a compact Lie group, four dimensional manifolds [3] or nilmanifolds [5]. If (M, ω) is aspherical we get a stronger statement.

The group of Hamiltonian diffeomorphisms of a closed symplectic manifold has the property that the evaluation map induces the trivial homomorphism of the fundamental group. The proof of this fact is nontrivial and can be found in McDuff-Salamon [4, Corollary 9.1.2]. Thus we can apply Theorem 2.1 in the Hamiltonian case and Corollary 2.2 gives the following result.

Theorem 3.1. *Let $(M, \omega) \rightarrow E \rightarrow B$ be a Hamiltonian fibration. If M is aspherical then the fibration is homotopy trivial. In particular, the Lalonde-McDuff conjecture holds for aspherical manifolds. $\square$*

Combining our argument with some known results we obtain the following more general theorem.

Theorem 3.2. *Let (M, ω) be a symplectic manifold. If the cohomology ring $H^*(M; \mathbb{Q})$ is generated by $H^2(M)$ and the Chern classes and the $\pi_1(M)$ -classes then the Lalonde-McDuff conjecture holds for (M, ω) .*

Proof. The fact that the second cohomology is contained in the image of the map induced by the inclusion of the fibre is proven in Lalonde-McDuff [3, Theorem 1.1].

The Chern classes are in the image of $i^* : H^*(E) \rightarrow H^*(M)$ because the tangent bundle TM is the pull-back of the bundle tangent to the fibers of $E \rightarrow B$.

Now the statement follows from Theorem 2.1 and the Leray-Hirsch theorem. $\square$

REFERENCES

- [1] D. H. Gottlieb. A certain subgroup of the fundamental group. *Amer. J. Math.*, 87:840–856, 1965.

- [2] Allen Hatcher. *Algebraic topology*. Cambridge University Press, Cambridge, 2002.
- [3] François Lalonde and Dusa McDuff. Symplectic structures on fiber bundles. *Topology*, 42(2):309–347, 2003.
- [4] Dusa McDuff and Dietmar Salamon. *J-holomorphic curves and symplectic topology*, volume 52 of *American Mathematical Society Colloquium Publications*. American Mathematical Society, Providence, RI, 2004.
- [5] Zofia Stępień. The Lalonde-McDuff conjecture for nilmanifolds. *Differential Geometry and its Applications*, 26(3):267–272, 2008.

MATHEMATICAL SCIENCES, UNIVERSITY OF ABERDEEN, MESTON BUILDING, ABERDEEN, AB243UE, SCOTLAND, UK, INSTITUTE OF MATHEMATICS, UNIVERSITY OF SZCZECIN, WIELKOPOLSKA 15, 70-451 SZCZECIN, POLAND

E-mail address: kedra@maths.abdn.ac.uk

URL: <http://www.maths.abdn.ac.uk/~kedra>